

T.I. GAISIN

The maximum principle for manifolds over a local algebra (2).

Let A be a finite-dimensional local commutative algebra over R , $\dim_R A = n$. In this work we consider compact manifolds over A , and prove that the real part of an A -differentiable function is constant. Also we find estimates for the dimensions of some spaces of 1-form.

For the theory of manifolds over algebras we refer the reader to [1],[2],[3]. We assume that all manifolds and mappings under consideration are smooth.

Each function $G : A^m \rightarrow A$ can be written as

$$G = e_i \cdot G^i, i = 0, \dots, n-1,$$

where e_i is a real basis of A . If G is A -differentiable, one can choose e_i such that

$$G(X^1, \dots, X^m) = g(x^1, \dots, x^m) + \sum_{|p|=1}^n \frac{1}{p!} \cdot \frac{D^p g}{dx^p} \cdot (X - x)^p, \quad (1)$$

where $p = (p_1, p_2, \dots, p_m)$, $p! = p_1! \cdot p_2! \cdot \dots \cdot p_m!$, $|p| = p_1 + p_2 + \dots + p_m$, and

$$(X - x)^p = (X^1 - x^1)^{p_1} \cdot (X^2 - x^2)^{p_2} \cdot \dots \cdot (X^m - x^m)^{p_m},$$

$$\frac{D^p g}{dx^p} = \frac{\partial^{|p|} g}{(\partial x^1)^{p_1} (\partial x^2)^{p_2} \dots (\partial x^m)^{p_m}}, X^j = e_i \cdot x^{j,i}, i = 1..n-1,$$

$$e_0 = 1 \in A, (e_k)^S = 0, k = 1..n-1, S \text{ is a positive integer,}$$

i.e. e_k belong to the radical $Rd(A)$ and there exists a pseudobasis e_l such that $l = 1..r, r \leq n-1$, $e_k = e_1^{s_1} \cdot \dots \cdot e_r^{s_r}$ (see [1],[2],[3]). We will call this basis the standard basis.

We will use the following proposition (see [1]).

Proposition.[1] Suppose that $\dim_R A = n$. Then the standard basis of A satisfies: $e_0 = 1 \in A$, $(e_1)^S = 0$, $(e_k)^S = 0$, and $k = 2..n-1$, where e_k is a real basis of an ideal in A .

Recall also that on a manifold M over A carries a foliation $\mathcal{F}$ called the canonical foliation.

1. Main theorem

Theorem 1. Let M be a manifold over A , $\dim_R A = n$, $\dim_A M = m$. Let g be the real part of an A -differentiable function on M .

Then for any leaf L of $\mathcal{F}$ there exists a leaf K of $\mathcal{F}$ such that $K \subset \overline{L}$ and $d(g)|_K = 0$.

Proof. By definition, we have

$$e_1 \cdot e_k = e_i \cdot v_k^i, \text{ where } i, k = 2..n-1, v_k^i \in R.$$

Evidently, $(e_1)^S = 0$ implies $e_1 \cdot e_1 = e_i \cdot v^i$. Then from (1) it follows that $G(X^1, \dots, X^m) = g(x^1, \dots, x^m) + e_i \cdot G^i = g + e_i \cdot g^i$. Here g^i are the coordinates of G with respect to a basis of A .

Let $\mathcal{A}$ be the maximal A -differentiable atlas. In the general case, with respect to a chart of $\mathcal{A}$, we have

$$g^1 = \frac{\partial g}{\partial x^j} \cdot x^{j,1} + b(x^1, \dots, x^m), \quad j = 1..m.$$

For the following reasoning it is important that g is a basic function of the canonical foliation on M and $b(x^1, \dots, x^m)$ depends only on the transversal coordinates $\{x^1, \dots, x^m\}$ of $\mathcal{F}$ (see [1],[2],[3],[4]).

Since M is compact, $\bar{L}$ is compact, too. Then g^1 has a minimum on $\bar{L}$ at a point $x \in K \subset \bar{L}$, where K is a leaf of $\mathcal{F}$. As is known, $K \subset \bar{L}$ [4]. One can take a chart (U, x^i) in $\mathcal{A}$ containing x such that x has coordinates $\{x^1, \dots, x^m, 0, \dots, 0\}$. If $d(g)|_K \neq 0$, then there exists a point $\tilde{x}$ in U with coordinates $\{x^1, \dots, x^m, x^{1,1}, \dots, x^{m,1}, 0, \dots, 0\}$ such that $g^1(x) > g^1(\tilde{x})$. Since $\tilde{x} \in K$ and x is a minimum point of g^1 , we obtain a contradiction. $\square$

From Theorem 1 it follows that $g^1|_{\bar{L}}$ achieves maximum and minimum on $\bar{L}/L$.

Theorem 2. *Let M be a compact manifold over A , ($\dim_R A = n$, $\dim_A M = m$). Suppose that e_i is a standard basis A . Let G be an A -differentiable function on M . Then*

- 1) *The real part g of G is constant;*
- 2) *The e_1 -component of G is constant on leaves of $\mathcal{F}$.*

Proof. From Theorem 1 it follows that all values of the basic function g are critical. Then the theorem assertion follows from the Sard Theorem. $\square$

Let us denote by $\tilde{e}_k$ the element of the standard basis of A such that $\tilde{e}_k \cdot e_l = 0$, where $l = 1..n-1$. This means that $\tilde{e}_k \cdot Rd(A) = 0$. The other elements of the standard basis of A we will denote by $\tilde{e}_j$.

Corollary 1. *Let M — be a compact manifold over A , and G be an A -differentiable function on M . Then*

$$G = a + f^k \cdot \tilde{e}_k,$$

where $a \in A$, and f^k are basic functions.

Proof. The corollary follows from Theorem 2 and the properties of the standard basis $\{e_i\}$. $\square$

2. Dimensions of 1-form spaces on M .

Let M be a manifold over A , Ω be the space of differentiable forms on M . Recall that an $\omega \in \Omega$ is said to be A -differentiable if ω and $d\omega$ are A -linear. Let us denote by Ω_A the space of A -differentiable forms on M . Note that

$$\Omega_A \subset \Omega_{A\text{-lin}} \subset A \otimes \Omega,$$

where $\Omega_{A\text{-lin}}$ is the space of A -linear forms[2].

From this we get the following short exact sequence of complexes:

$$0 \rightarrow \Omega_A \rightarrow A \otimes \Omega \rightarrow \frac{A \otimes \Omega}{\Omega_A} \rightarrow 0.$$

The left complex (Ω_A, d) is called the complex of A -differentiable forms. Then we obtain the cohomology exact sequence

$$0 \rightarrow H_A^0 \rightarrow A \otimes H^0 \rightarrow H_{/A}^0 \rightarrow H_A^1 \rightarrow A \otimes H^1 \rightarrow \dots,$$

where the groups without lower indices are the de Rham cohomology groups of M , H_A^* are the cohomology groups of (Ω_A, d) , and $H_{/A}^*$ are the cohomology groups of the quotient complex $(\Omega_{/A}, d)$.

Lemma 1. $\dots \rightarrow 0 \rightarrow H_A^1 \rightarrow A \otimes H^1 \rightarrow \dots$ is a monomorphism.

For every j_0 , let

$$\check{\Omega}_{AR}^{s,j_0} = \{\check{\omega}^{j_0} \in \Omega^s \mid \exists \omega \in \Omega_A, \omega = \check{e}_{j_1} \cdot \check{\omega}^{j_1} + \check{e}_{j_0} \cdot \check{\omega}^{j_0} + \check{e}_{j_2} \cdot \check{\omega}^{j_2} + \check{e}_k \cdot \check{\omega}^k\}.$$

Note that $\check{\Omega}_{AR}^{s,j_0} \subset \Omega^s$.

For each j_0 , let

$$Z\check{\Omega}_{AR}^{s,j_0} = \{\check{\omega}^{j_0} \in \Omega^s \mid \exists \omega \in (\Omega_A \cap \ker\{d\}), \omega = \check{e}_{j_1} \cdot \check{\omega}^{j_1} + \check{e}_{j_0} \cdot \check{\omega}^{j_0} + \check{e}_{j_2} \cdot \check{\omega}^{j_2} + \check{e}_k \cdot \check{\omega}^k\}$$

Note that $Z\check{\Omega}_{AR}^{s,j_0} \subset \check{\Omega}_{AR}^{s,j_0}$.

Theorem 3. *Let M be a compact manifold over A , $\dim_R A = n$, and $\dim_A M = m$. Then the spaces $Z\check{\Omega}_{AR}^{1,j}$ have the finite dimension and*

$$\dim\{Z\check{\Omega}_{AR}^{1,j}\} \leq \dim\{A \otimes H^1(M)\}.$$

Proof. Let $\omega, \varsigma \in [\omega] \in H_A^1$, and $\omega \in \Omega_A$, $\omega = e_i \cdot \omega^i = \check{e}_j \cdot \check{\omega}^j + \check{e}_k \cdot \check{\omega}^k$, $\varsigma \in \Omega_A$, $\varsigma = e_i \cdot \varsigma^i = \check{e}_j \cdot \check{\varsigma}^j + \check{e}_k \cdot \check{\varsigma}^k$. From Corollary 1 it follows that $\check{\Omega}_{AR}^{0,j} \cong R$. Hence $\check{\omega}^j = \check{\varsigma}^j$ and $\dim\{H^1(M)\} < \infty$.

The second statement follows from Lemma 1. $\square$

References.

[1] V.V. Vishnevskii, A.P. Shirokov, V.V. Shurygin. *Spaces over algebras*, Kazan University Press.1985.

[2] Shurygin, V. V. *Jet bundles as manifolds over algebras*. (Russian) Translated in J. Soviet Math. 44 (1989), no. 2, 85–98. Itogi Nauki i Tekhniki, Problems of geometry, Vol. 19 (Russian), 3–22, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1987. (MR0933562 (89d:58006)).

[3] Shurygin, V. V. *Manifolds over algebras and their application in the geometry of jet bundles*. (Russian) Uspekhi Mat. Nauk 48 (1993), no. 2(290), 75–106; translation in Russian Math. Surveys 48 (1993), no. 2, 75–104 (MR1239861 (94i:58004)).

[4] Molino P. *Riemannian foliations*. – Birkhäuser, 1988.

N. G. CHEBOTAREV RESEARCH INSTITUTE OF MATHEMATICS AND MECHANICS, GEOMETRY DEPARTMENT, UNIVERSITETSKAYA, 17, KAZAN, TATARSTAN: 420008, RUSSIAN FEDERATION

E-mail address: Tagir.Gaisin@ksu.ru